

The finite temperature transition for 3-flavour lattice QCD at finite isospin density*

J. B. Kogut^{a†} and D. K. Sinclair, ^{b‡}

^aDepartment of Physics, University of Illinois, 1110 West Green Street, Urbana, IL 61801, USA

^bHEP Division, Argonne National Laboratory, 9700 South Cass Avenue, Argonne, IL 60439, USA

We simulate 3-flavour lattice QCD with a small chemical potential μ_I for isospin, at temperatures close to the finite temperature transition. Using quark masses just above the critical mass for zero chemical potential, we determine the position of the transition from hadronic matter to a quark-gluon plasma as a function of μ_I . We see evidence for a critical endpoint where the transition changes from a crossover to a first-order transition as μ_I is increased. We argue that QCD at finite μ_I and QCD at finite quark-number chemical potential μ should behave similarly in this region.

1. Introduction

QCD at finite temperature and quark-number chemical potential, μ , is of relevance to the physics of relativistic heavy-ion collisions. Different parts of the phase diagram are accessible to the various heavy-ion accelerators — RHIC, CERN heavy-ion, AGS and SIS.

For the region of small μ and temperatures close to the finite temperature transition, various techniques have been developed to circumvent the problems associated with the complex fermion determinant at finite μ [1,2,3,4]. We have chosen to study the finite temperature transition with a finite chemical potential μ_I for isospin, which is equivalent to simulating QCD at finite μ with $2\mu = \mu_I$ using only the magnitude of the determinant and ignoring the phase. Because the fermion phase is relatively well behaved at small μ , it can be argued that the positions of the transitions and probably their nature with and without this phase, should be the same.

For 2 flavours we found no sign of a critical endpoint for $\mu_I < m_\pi$, the maximal range over which finite μ_I and finite μ transitions could be expected to be related [5]. Here we describe our simulations for 3 flavours. Not only is this more physical, but

for $m < m_c(0)$, the finite temperature transition is first order at $\mu_I = 0$. At $m = m_c(0)$ it becomes second-order, in the Ising universality class. For $m > m_c(0)$ the $\mu_I = 0$ transition is a crossover. It is expected that the critical endpoint moves continuously from $\mu_I = 0$ for $m = m_c(0)$ to finite μ_I , as m is increased. Thus we should be able to keep this endpoint as close to $\mu_I = 0$ as we please, by choosing m to lie just above $m_c(0)$. The preliminary results of our simulations at such a mass, reported here, do show such a critical endpoint at a modest value of μ_I .

In section 2 we describe our simulations and report preliminary results. Our conclusions are given in section 3.

2. 3-flavour QCD at finite μ_I and T

The staggered fermion part of our lattice action is

$$S_f = \sum_{sites} \left[\bar{\chi} [\not{D}(\frac{1}{2}\tau_3\mu_I) + m]\chi + i\lambda\epsilon\bar{\chi}\tau_2\chi \right]. \quad (1)$$

For these simulations, which are at $\mu_I < m_\pi$, we set the symmetry breaking parameter $\lambda = 0$.

We are performing simulations on $8^3 \times 4$ and $12^3 \times 4$ lattices at $m = 0.03, 0.035$ and 0.04 . [For $N_t = 4$, $m_c(0) = 0.0331(12)$ [6].] We use standard hybrid molecular-dynamics simulations. Since with such small lattices one can get 2-state

*Talk presented by D. K. Sinclair, Lattice2004, Fermilab.

†Supported in part by NSF grant NSF PHY03-04252.

‡Supported by DOE contract W-31-109-ENG-38.

signals even for crossovers, we use fourth-order Binder cumulants (B_4) [7] for $\bar{\psi}\psi$ to get quantitative information about the transition:

$$B_4(X) = \frac{\langle (X - \langle X \rangle)^4 \rangle}{\langle (X - \langle X \rangle)^2 \rangle^2}, \quad (2)$$

where X is any observable. Since we use 5 noise vectors for each configuration to estimate $\bar{\psi}\psi$ (and the number densities), we are able to obtain an unbiased estimate of $B_4(\bar{\psi}\psi)$, as well as the susceptibility $\chi_{\bar{\psi}\psi}$.

$$\chi_X = V \langle X^2 - \langle X \rangle^2 \rangle. \quad (3)$$

The β of the transition is determined by the positions of the maxima of the various susceptibilities, or from the minimum of $B_4(\bar{\psi}\psi)$. (We have checked the consistency of these two methods.)




Figure 1. β_c as functions of μ_I^2 from minima of B_4 . The bottom point is for $m = 0.03$. The middle set of points and line are for $m = 0.035$. The top set of points and line are for $m = 0.04$.

Figure 1 shows the transition β s, β_c , as functions of μ_I for the 3 quark masses. The fits shown

are

$$\begin{aligned} m = 0.035 & : \quad \beta_c = 5.1489(5) - 0.145(6)\mu_I^2 \\ m = 0.040 & : \quad \beta_c = 5.1569(3) - 0.147(4)\mu_I^2 \end{aligned} \quad (4)$$

which reflect the expected analyticity in μ_I^2 . β_c was obtained in each case from the position of the minimum of $B_4(\bar{\psi}\psi)$. Ferrenberg-Swendsen reweighting was used to interpolate between the β s of our simulations. These positions were in good agreement with those obtained from the maxima of the corresponding susceptibilities.




Figure 2. Binder cumulants for $m = 0.035$ as functions of μ_I . The dashed line is at $B_4 = 1.604$, the 3-d Ising value.

For a lattice of infinite volume, the Binder cumulant $B_4(\bar{\psi}\psi)$ is 3 for a crossover and 1 for a first-order transition. The critical endpoint is expected to be in the universality class of the 3-d Ising model, for which $B_4(\bar{\psi}\psi) = 1.604(1)$. While B_4 for a crossover or a first-order transition approaches its infinite volume limit relatively slowly with volume, the approach for a second-order phase transition is fast and approaches the limiting value on relatively small lattices [8]. Figure 2

shows the Binder cumulant at the transition for our runs at $m = 0.035$ with $\mu_I = 0$ and $\mu_I = 0.2$ for both $8^3 \times 4$ and $12^3 \times 4$ lattices. Despite the sizable error bars, these are the results of runs of 160,000 trajectories at the β closest to the transition.

B_4 at $\mu_I = 0$ lies above the Ising value while that at $\mu_I = 0.2$ lies below this value for both lattice sizes. Straight lines through these 2 points for the two lattice sizes cross at some μ_I with $0.0 < \mu_I < 0.2$, and at a B_4 value close to the Ising value. These preliminary results would indicate that there is a critical endpoint at $\mu_I = \mu_{IE}$ with $0.0165 \lesssim \mu_{IE}^2 \lesssim 0.0231$.

For the m, μ_I values at which we have run and for β s in the neighbourhood of the transition on an $8^3 \times 4$ lattice we estimate that the phase of the fermion determinant θ obeys,

$$\langle \theta^2 \rangle \approx (5-12)\mu_I^2. \quad (5)$$

We have argued in [5] that $\langle \cos \theta \rangle > 0.5$ is a reasonable estimate of the range over which one can neglect the phase of the fermion determinant in determining the position and nature of the finite temperature transition [5]. If so, the finite μ_I and μ behaviour of this transition should agree for $\mu_I \leq 0.25 - 0.45$, which includes our estimate for the critical endpoint for $m = 0.035$.

Using $m_c(0) = 0.0331$ we then predict that

$$\frac{m(\mu_I)}{m(0)} \approx 1 + 7 \left(\frac{\mu_I}{2\pi T_c} \right)^2 \quad (6)$$

3. Conclusions

We are simulating 3-flavour lattice QCD at finite μ_I and temperatures close to the finite temperature transition.

We observe the slow decrease of β and hence temperature, of the finite temperature transition with μ_I . Arguing that, in this range of μ_I , the dependence on μ_I and 2μ should be the same we find a falloff about 20% slower than that observed by de Forcrand and Philipsen [3] At least some of the difference is due to omitting $\mathcal{O}(\mu_I^4)$ terms.

We have obtained preliminary evidence for a critical endpoint for $m = 0.035$ (just above $m_c(0)$) from simulations at $\mu_I \leq 0.2$. However,

simulations on an $8^3 \times 4$ lattice at $\mu_I = 0.375$ yield $B_4 = 1.64(2)$, which does not support this interpretation. This is probably because 0.375 is too close to $\mu_c = m_\pi$, but this needs to be clarified. We have seen no evidence for a critical endpoint at $m = 0.04$. Whether this is consistent with what we see at $m = 0.03$ should also be checked.

Our prediction for the μ_I dependence of $m_c(\mu_I)$ appears inconsistent with both de Forcrand and Philipsen [3] and the Bielefeld-Swansea collaboration [2], lying in between their inconsistent results.

Clearly much more needs to be done.

Acknowledgements

DKS would like to thank Philippe de Forcrand for helpful discussions on Binder cumulants. These simulations were performed on the LCRC Jazz cluster at Argonne, the Tungsten cluster at NCSA and local PCs.

REFERENCES

1. Z. Fodor and S. D. Katz, Phys. Lett. B **534**, 87 (2002) [arXiv:hep-lat/0104001]; JHEP **0203**, 014 (2002) [arXiv:hep-lat/0106002]; JHEP **0404**, 050 (2004) [arXiv:hep-lat/0402006].
2. C. R. Allton *et al.*, Phys. Rev. D **66**, 074507 (2002) [arXiv:hep-lat/0204010]; F. Karsch, *et al.*, Nucl. Phys. Proc. Suppl. **129**, 614 (2004) [arXiv:hep-lat/0309116].
3. P. de Forcrand and O. Philipsen, Nucl. Phys. B **642**, 290 (2002) [arXiv:hep-lat/0205016]; Nucl. Phys. B **673**, 170 (2003) [arXiv:hep-lat/0307020].
4. M. D'Elia and M. P. Lombardo, Phys. Rev. D **67**, 014505 (2003) [arXiv:hep-lat/0209146].
5. J. B. Kogut and D. K. Sinclair, arXiv:hep-lat/0407027.
6. F. Karsch, E. Laermann and A. Peikert, Nucl. Phys. B **605**, 579 (2001) [arXiv:hep-lat/0012023].
7. K. Binder, Z. Phys. B **43**, 119 (1981).
8. F. Karsch, E. Laermann and C. Schmidt, Phys. Lett. B **520**, 41 (2001) [arXiv:hep-lat/0107020].